

LECTURE NOTES OF

# **ENGINEERING MATHEMATICS-III**

3<sup>RD</sup> SEMESTER ETC



**PREPARED BY- ROLEX MAHARANA**

**GOVERNMENT POLYTECHNIC DHENKANAL**

**DEPARTMENT OF ELECTRONICS AND TELECOMMUNICATION ENGINEERING**

## Chapter - 1

## Matrices

### Basic concepts of matrices:-

- A matrix is a rectangular arrangement (or) of numbers either real or complex on both.
- A matrix with  $m$  rows &  $n$  columns is called  $m \times n$  matrix or  $m$  by  $n$ .
- A rectangular arrangement of elements of the form

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

( $m$ ) one rows & ( $n$ ) one columns.

we can also write it as  $[a_{ij}]_{m \times n}$  or  $(a_{ij})_{m \times n}$

Ex

$$\begin{bmatrix} 1 & 5 & 7 \\ 2 & 8 & 9 \\ 4 & 2 & 7 \end{bmatrix}_{3 \times 3} \quad \begin{bmatrix} 5 & 7 & 4 \\ 2 & 1 & 0 \end{bmatrix}$$

### Different types of matrices:-

#### Row Matrix:-

A matrix having a single row is called row matrix.

→ It is in the form  $[a_{ij}]_{1 \times n}$  ( $a_{11} a_{12} a_{13}$ )

Ex  $[123]_{1 \times 3}$

#### Column Matrix:-

A matrix having a single column is called column matrix.

In the form  $[a_{ij}]_{m \times 1} = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}_{3 \times 1}$

## Chapter - 1

## Matrices

### Basic concepts of matrices:-

- A matrix is a rectangular arrangement (or) of numbers either real or complex or both.
- A matrix with  $m$  rows &  $n$  columns is called  $m \times n$  matrix or  $m$  by  $n$ .
- A rectangular arrange of elements of the form

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

( $m$ ) one rows & ( $n$ ) one columns.

we can also write it as  $[a_{ij}]_{m \times n}$  or  $(a_{ij})_{m \times n}$

Ex

$$\begin{bmatrix} 1 & 5 & 7 \\ 2 & 8 & 9 \\ 4 & 2 & 7 \end{bmatrix}_{3 \times 3} \quad \begin{bmatrix} 5 & 7 & 4 \\ 2 & 1 & 0 \end{bmatrix}$$

### Different types of matrices:-

#### Row Matrix:-

A matrix having a single row is called row matrix.

→ It is in the form  $[a_{ij}]_{1 \times n}$  ( $a_{11} a_{12} a_{13}$ )

Ex  $[123]_{1 \times 3}$

#### Column Matrix:-

A matrix having a single column is called column matrix.

In the form  $[a_{ij}]_{m \times 1} = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}_{3 \times 1}$

### Rectangular Matrix:-

A matrix of order  $m \times n$  is said to be a rectangular matrix if  $m \neq n$ .

### Square Matrix:-

#### Note:-

The element of square matrix  $A = [a_{ij}]_{m \times n}$  are classified into 3 types.

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}_{n \times 1}$$

(1) The element  $a_{11}, a_{22}, a_{33}, \dots, a_{nn}$  are called diagonal element.

(2) The elements  $a_{ij}$   $i < j$  are upper diagonal elements.

(3) The elements  $a_{ij}$ ,  $i > j$  are lower diagonal element.

### Diagonal Matrix:-

$$\begin{bmatrix} 5 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 8 \end{bmatrix}$$

### Scalar Matrix:-

A matrix is said to be a scalar matrix if all the diagonal elements are same.

Ex

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$



### Null Matrix :-

A matrix said to be a null matrix if all the entries are zero, is zero matrix denoted by zero.

### Unit Matrix and Identity Matrix :-

A matrix said to be a unit matrix if all the entries in the leading diagonal if  $a_{ij} = \begin{cases} 0, & i \neq j \\ 1, & i = j \end{cases}$

Ex

$$I_1 = [1]_{1 \times 1}$$

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}_{2 \times 2}$$

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3}$$

### Addition of matrix :-

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Ex  $\begin{bmatrix} 2 & 0 & 0 \\ 1 & 0 & 0 \\ 5 & 4 & 7 \end{bmatrix} \rightarrow \text{Lower}$

### Lower Triangular Matrix :-

- Lower triangular matrix can have non-zero entries only on and below the main diagonal.
- Any entries on the main diagonal on the triangular matrix may be zero or not other wise a square matrix if  $[a_{ij}] = 0, i < j$ , that is element above the leading diagonal

or if all its upper diagonal is zero.

### Upper Triangular Matrix :-

Upper triangular matrix are square matrices that can have non-zero matrices. Only on above the main diagonal. where, as any entry below the diagonal must be zero otherwise a square matrix  $[a_{ij}]$  is called an upper triangular matrix. if  $a_{ij} =$

that is element below the leading diagonal are zero or if all its lower diagonal element zero.

ex

$$\begin{bmatrix} 2 & 0 & 2 \\ 0 & 7 & 5 \\ 0 & 0 & 0 \end{bmatrix}$$

### Addition of Matrix :-

Let  $A = [a_{ij}]$  be a matrix of order  $(m \times n)$  and

$B = [b_{ij}] m \times n$  then their addition before

than

$$A+B = [a_{ij} + b_{ij}] m \times n$$

where  $i = 1, 2, \dots, m$

$j = 1, 2, 3, \dots, n$

That is addition of two matrices of same order is often by adding the element in the corresponding position.

ex

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 5 & 2 & 7 \end{bmatrix}_{2 \times 3}$$

$$B = \begin{bmatrix} 6 & 1 & 2 \\ 2 & 3 & 4 \end{bmatrix}_{2 \times 3}$$

$$A+B = \begin{bmatrix} 1+6 & 2+1 & 3+2 \\ 5+2 & 3+2 & 7+4 \end{bmatrix}_{2 \times 3}$$

$$= \begin{bmatrix} 7 & 3 & 5 \\ 7 & 5 & 11 \end{bmatrix}_{2 \times 3}$$

### Subtraction of Matrix :-

The subtraction of two matrix A & B of same order is defined as,

$$\boxed{[A-B] = (A+(-B))}$$

where  $(-B)$  is the negative (-ve) matrix of the  $[B]$

### Transpose Matrices :-

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}_{2 \times 2} = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}_{2 \times 2}$$

$$A = \begin{bmatrix} 1 & 3 & 7 \\ 2 & 1 & 9 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 7 & 9 \end{bmatrix}_{3 \times 2}$$

### Symmetric Matrix :-

$$A = \begin{bmatrix} 2 & 4 \\ 4 & 3 \end{bmatrix}_{2 \times 2}$$

$$A^T = \begin{bmatrix} 2 & 4 \\ 4 & 3 \end{bmatrix}_{2 \times 2}$$

or

$$A = \begin{bmatrix} 0 & 1 & -3 \\ -1 & 0 & -2 \\ -3 & -2 & 0 \end{bmatrix}_{3 \times 3}$$

$$-A = \begin{bmatrix} 0 & -1 & +3 \\ +1 & 0 & +2 \\ -3 & +2 & 0 \end{bmatrix}_{3 \times 3}$$

$$A^T = \begin{bmatrix} 0 & -1 & 3 \\ 1 & 0 & -2 \\ -3 & -2 & 0 \end{bmatrix}_{3 \times 3}$$

## Matrix multiplication:-

Let  $A = [a_{ij}]$  be a matrix of order  $m \times n$  and  $B = [b_{jk}]$  matrix of order  $n \times p$  so that, number of columns in  $A$  = number of rows in  $B$ .

Then the product  $A \cdot B$  is well defined and it will be matrix of order  $m \times p$ , whose elements are given by  $c_{ik} = \sum_{j=1}^n a_{ij} b_{jk}$

$b_{1k} + a_{i2} b_{2k} + \dots + a_{in} b_{nk}$  ( $i, k$ ) element of  $A \cdot B$ .

Ex 1

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3}$$

$$B = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}_{3 \times 3}$$

$$AB = a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31} \quad a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} \quad a_{11}b_{13} + a_{12}b_{23} + a_{13}b_{33}$$

$$AB = a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31}$$



Properties of matrix multiplication:-

For any 3 matrices  $A, B, C$  conformable for multiplication and Scalar.

$$(1) K(AB) = (KA)B = A(KB)$$

$$(2) A(BC) = (AB)C \text{ (Associative)}$$

$$(3) (A+B)C = AC + BC$$

$$C(A+B) = CA + CB$$

Sub matrix & minors:-

\* Any matrix often by omitting some rows or columns or both of a given  $m \times n$  matrix ' $A$ ' is called a sub-matrix of ' $A$ '. Thus  $\begin{bmatrix} a_1 & b_1 \\ a_2 & b_2 \end{bmatrix}$  is a sub-matrix of  $\begin{matrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{matrix}$

Rank of a matrix:-

A number ' $r$ ' is said to be a ~~matrix~~ rank of a non-zero  $n \times m$  matrix.

if (i) there is at least one  $(r \times r)$  square matrix of  $A$  whose determinant is not equal to zero.

\* (ii) The determinant of every  $(r+1)$  rowed square sub-matrix in  $A$  is zero.

How do find rank of matrix:-

To find the rank of matrix, we will transform that matrix into its echelon form in linear algebra matrix a row echelon form if it has the say resulting from a Gaussian elimination all rows consisting of only zero are at the bottom. The leading coefficient (also called pivot of non zero - rows is always strictly to the right of the leading coefficient of the row above it.) then determine the rank of number non-zero rows.



$$A = \begin{bmatrix} 2 & 4 & 6 \\ 4 & 8 & 12 \end{bmatrix}_{2 \times 2}$$

condition :-

\* If a matrix is of order  $m \times n$  how  $\leq 0$   $P(A) \leq \min(m, n)$   
 = minimum of  $(m, n)$ .

\* If  $A$  is order  $m \times n$  an determinant  $|A| \neq 0$  then the rank of  $A = n$

\* If  $A$  is of order  $m \times n$  &  $|A| = 0$ , then the rank of  $A$  will be less than  $n$ .

Ex

find the Rank of

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}_{3 \times 3}$$

Ans

$$\text{Let } A = \begin{bmatrix} \oplus & \ominus & \oplus \\ 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}_{3 \times 3}$$

$$\text{where } P(A) \leq \min(3, 3)$$

$$= 1(15-16) - 2(10-12) + 3(8-9)$$

$$= (15-16) - 2(-2) + 3(-1)$$

$$= -1 + 4 - 3$$

$$= -4 + 4$$

$$= 0$$

Here  $A$  is a singular square matrix in which there is at least one  $(2 \times 2)$  sub matrix.

For example

$$A_1 = \begin{bmatrix} 3 & 4 \\ 4 & 5 \end{bmatrix}_{2 \times 2}$$

$$15 - 16 = -1 \neq 0$$

Hence, The rank of  $A$  is 2. which is less than the order of 3 of singular square matrix.

Linear system of eq's :- (Existence, uniqueness)

A linear eq<sup>n</sup> with 'n' unknowns  $x_1, x_2, \dots, x_n$  is an eq<sup>n</sup> is of the form  $a_1x_1 + a_2x_2 + \dots + a_nx_n = b$  — (1)

In the above eq<sup>n</sup> if  $b = 0$  then it is called homogeneous linear eq<sup>n</sup>.

In contrast eq<sup>n</sup> (1) is a non homogeneous linear eq<sup>n</sup>.

Consider a set of 'M' non homogeneous eq<sup>n</sup> with 'n'.

$$\left. \begin{array}{l} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + \dots + a_{mn}x_n = b_m \end{array} \right\} \text{--- (2)}$$

\* At least one set of values of  $x_1, x_2, x_3, \dots, x_n$  can be found satisfying all the eq's then the set of eq's are known as consistent.

If no such set exist then the eq's are said to be inconsistent.

Ex  $\rightarrow$  (i)  $2x + 5y = 9$   
 $x - y = 1$

(ii)  $x + 2y = 7$   
 $4x + 8y = 28$

(iii)  $2x + 3y = 5$   
 $4x + 6y = -8$

(i)  $2x + 5y = 9$

$2x(x - y = 1)$

$$\begin{array}{r} 2x + 5y = 9 \\ 2x - 2y = 2 \\ \hline (-) \quad (+) \quad (-) \end{array}$$

$7y = 7$

$\Rightarrow y = 1$

(Consistent)

$$\textcircled{2} \quad x + 2y = 7$$

$$4x + 8y = 28$$

$$\Rightarrow x = 7 - 2y$$

$\Rightarrow$  no. of sol<sup>n</sup>s

$\Rightarrow$  consistent

$$\textcircled{3} \quad 2 \times (2x + 3y = 5)$$

$$4x + 6y = -8$$

$$\begin{array}{r} 4x + 6y = 10 \\ 4x + 6y = -8 \\ \hline 0 = 18 \end{array}$$

$\Rightarrow$  This eq<sup>n</sup> have no sol<sup>n</sup>

$\Rightarrow$  Inconsistent.

The association of homogenous is given by

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0$$

$\vdots$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0$$

—  $\textcircled{3}$

The coefficient matrix  $A$  is given by

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} = A$$

is called coefficient matrix denoted by  $A$ .

Then 
$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{bmatrix}$$

is called the augmented matrix which is denoted by  $A_b$  or  $(A, b)$

The system of non-homogeneous eq<sup>n</sup>s (2) may be put in the form

$$AX = B$$

where  $x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$   $\neq$   $B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$

consistency of system of linear eq<sup>n</sup> :-

consider the following m eq<sup>n</sup> n unknown

$$\left. \begin{array}{l} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{array} \right\} \text{--- (1)}$$

which is in the matrix form

$$AX = B$$

where

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$



$$X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \& \quad B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

Here,  $A$  is called co-efficient matrix

$B$  is called right hand side matrix  
with the help of  $A$  and  $B$  consider

$$K = [A/B]$$

$$\Rightarrow A = \left[ \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right]$$

which is known as augmented matrix.

A sol<sup>n</sup> of eq<sup>n</sup> ① defined by set of values of the -  
variable  $x_1, x_2, \dots, x_n$  which satisfy.

If the system given by eq<sup>n</sup> ① has a sol<sup>n</sup>, it is  
called consistent system. otherwise this is called ~~in~~-  
inconsistent system.

In fact a consistent system has either unique sol<sup>n</sup> or  
~~infinite~~ infinitely many sol<sup>n</sup>.

Rouche's theorem :-

The system of equation one if was only co-efficient matrix  $A$  and the augmented matrix  $K$  are of some rank otherwise the system is inconsistent.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$K = \left[ \begin{array}{ccc|c} a_{11} & a_{12} & a_{13} & b_1 \\ a_{21} & a_{22} & a_{23} & b_2 \\ a_{31} & a_{32} & a_{33} & b_3 \end{array} \right]$$

$$\begin{aligned} \star \text{ Rank of } A &= \text{Rank of } K \\ &= r \quad (r \leq \text{the smaller of } m \text{ \& } n) \end{aligned}$$

The equation (1) can by suitable row operations be reduce to

$$b_{11}x_1 + b_{12}x_2 + \dots + b_{1n}x_n = k_1$$

And the remaining  $(m-r)$  equation being all of the form

$0x_1 + 0x_2 + \dots + 0x_n = 0$

\* The equation (2) will have a solution through  $n-r$  of the unknown may be chosen arbitrarily. The solution will be unique when  $r=n$ .

Hence the equations (i) are consistent.

\* Rank of 'A' < rank of K. In particular, let the rank of K be  $r+1$ . In this case the equations (i) will reduce, by suitable row operations to.

$$b_{11}x_1 + b_{12}x_2 + \dots + b_{1n}x_n = k_1$$

$$0x_1 + b_{22}x_2 + \dots + b_{2n}x_n = k_2$$

⋮

⋮

⋮

⋮

⋮

$$0x_1 + 0x_2 + \dots + b_{rn}x_n = k_r$$

$$0x_1 + 0x_2 + \dots + 0x_n = k_{r+1}$$

and remaining  $m-(r+1)$  equations are of the form  $0x_1 + 0x_2 + \dots + 0x_n = 0$ . Clearly the  $(r+1)$ th equation can not be satisfied by any set of values for the unknowns. Hence the equations (i) are inconsistent.

Ex ① solve  $x + 2y - z = 3$   
 $3x - y + 2z = 1$   
 $2x - 2y + 3z = 2$

Ans

Given eqns are

$$\left. \begin{array}{l} x + 2y - z = 3 \\ 3x - y + 2z = 1 \\ 2x - 2y + 3z = 2 \end{array} \right\} \text{--- ①}$$

which can write in the form

$$AX = B$$

$$\begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

where

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 3 \end{bmatrix} \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad B = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

The given eq<sup>n</sup> which can be represented in augmented matrix form

$$K = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \\ 2 & -2 & 3 & 2 \end{bmatrix}$$

$$R_2 = R_2 - 3R_1$$

$\sim$  = equivalent symbol

$$\sim \begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & -7 & 5 & -8 \\ 2 & -2 & 3 & 2 \end{bmatrix}$$

$$R_3 = R_3 - 2R_1$$

$$\sim \begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & -7 & 5 & -8 \\ 0 & -6 & 5 & -4 \end{bmatrix}$$

$$(C_2 \leftrightarrow C_3)$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 3 \\ 0 & 5 & -7 & -8 \\ 0 & 5 & -6 & -4 \end{bmatrix}$$

$$R_3 = R_3 - R_2$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 3 \\ 0 & 5 & -7 & -8 \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

$$\rho(K) = 3$$

$$\rho(A) = 3$$

Here  $\rho(K) = \rho(A) \Rightarrow$  system are consistent

Next to find the soln of eqn ①

$$x - y + 2z = 3$$

$$5y - 7z = -8$$

$$z = 4$$

Then to find y we can solve.

$$5y - 7z = -8$$

$$\Rightarrow 5y - 7 \times 4 = -8$$

$$\Rightarrow 5y - 28 = -8$$

$$\Rightarrow 5y = 28 - 8$$

$$\Rightarrow 5y = 20$$

$$\Rightarrow y = 20/5 = 4$$



Then,

$$x - y + 2z = 3$$

$$\Rightarrow x - 4 + 2 \times 4 = 3$$

$$\Rightarrow x - 4 + 8 = 3$$

$$\Rightarrow x + 4 = 3$$

$$\Rightarrow x = -1$$

Hence,  $x = -1$

$$y = 4$$

$z = 4$  are the required solution.

Q. Solve the following system completely.

$$2x - y + 3z = 3$$

$$x + 2y - z - 5w = 4$$

$$x + 3y - 2z - 7w = 5$$

Sol<sup>n</sup> writing the above eq<sup>n</sup> in matrix form  $AX = B$

we get

$$\begin{bmatrix} 2 & -1 & 3 & 0 \\ 1 & 2 & -1 & -5 \\ 1 & 3 & -2 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$$

where,

$$A = \begin{bmatrix} 2 & -1 & 3 & 0 \\ 1 & 2 & -1 & -5 \\ 1 & 3 & -2 & -7 \end{bmatrix} \quad X = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

$$B = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$$

Here the augmented matrix

$$K = \left[ \begin{array}{cccc|c} 2 & -1 & 3 & 0 & 3 \\ 1 & 2 & -1 & -5 & 4 \\ 1 & 3 & -2 & -7 & 5 \end{array} \right]$$

$$R_2 \leftrightarrow R_1$$

$$\sim \left[ \begin{array}{cccc|c} 1 & 2 & -1 & -5 & 4 \\ 2 & -1 & 3 & 0 & 3 \\ 1 & 3 & -2 & -7 & 5 \end{array} \right]$$

$$R_2 = R_2 - 2R_1 \text{ \& } R_3 = R_3 - R_1$$

$$\sim \left[ \begin{array}{cccc|c} 1 & 2 & -1 & -5 & 4 \\ 0 & -5 & 5 & 10 & -5 \\ 0 & 1 & -1 & -2 & 1 \end{array} \right]$$

$$R_2 = \frac{1}{5} R_2$$

$$\sim \left[ \begin{array}{cccc|c} 1 & 2 & -1 & -5 & 4 \\ 0 & -1 & 1 & 2 & -1 \\ 0 & 1 & -1 & -2 & 1 \end{array} \right]$$

$$R_3 = R_3 + R_2$$

$$\sim \left[ \begin{array}{cccc|c} 1 & 2 & -1 & -5 & 4 \\ 0 & -1 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\text{So } f(K) = 2$$

$$f(A) = 2$$

Hence

$$f(K) = f(A)$$

So, the equations are

$$x + 2y - z - 5w = 4 \quad \text{--- (1)}$$

$$-y + z + 2w = -1$$

$$\Rightarrow y - z - 2w = 1 \quad \text{--- (2)}$$

$$\text{Let } z = k_1 \text{ \& } w = k_2$$

So the eqn (2)

$$y = 1 + z + 2w$$

$$\text{in eqn (1)} = 1 + k_1 + 2k_2$$

$$x + 2(1 + k_1 + 2k_2) - (k_1 - 5k_2) = 4$$

$$\Rightarrow x + 2 + 2k_1 + 4k_2 - k_1 - 5k_2 = 4$$

$$\Rightarrow x + k_1 - k_2 = 2$$

$$\Rightarrow x = 2 + k_2 - k_1$$

Hence,

$$x = 2 + k_2 - k_1$$

$$y = 1 + z + 2w$$

$$z = k_1 \text{ \& } w = k_2$$

where  $k_1$  and  $k_2$  is constant.

## Chapter - 2      COMPLEX NUMBER

Real Number :-

Real numbers are numbers that includes both rational and irrational numbers. Rational numbers such as integer  $(-2, -1, 0, 1, 2 \text{ etc})$ , fractions  $(\frac{1}{2}, \frac{5}{7}, 2.5, 7.1)$  and irrational numbers such as  $(\sqrt{3}, \sqrt{5}, \sqrt{2}, \pi (\frac{22}{7}))$ .

\* Square of a positive real number is positive and that of a negative real is also positive. So, there is no real number - whose square is negative. So, we are to create a new kind of number. we define a square root of a negative number as imaginary number particularly  $\sqrt{-1} = i$  the basic imaginary number ~~particularly  $\sqrt{-1} = i$~~ . Then,  $\sqrt{-4} = 2i$ ,  $\sqrt{-9} = 3i$ .  $\sqrt{2} = \sqrt{2} i$  etc.

Taking  $i = \sqrt{-1}$

$$i^2 = \sqrt{-1} \times \sqrt{-1} = (\sqrt{-1})^2 = -1$$

$$i^3 = i^2 \times i = -i$$

$$i^4 = i^2 \times i^2 = 1$$

Since  $i^4 = 1$ ,

$$i = i^5 = i^9 = i^{13} + \dots \dots \dots i^{4n+1}$$

when  $n$  is integer.

$$i^2 = i^6 = i^{10} = \dots \dots \dots = i^{4n+2}$$

$$i^3 = i^7 = i^{11} = \dots \dots \dots = i^{4n+3}$$

$$i^4 = i^8 = i^{12} = i^{16} = \dots \dots \dots = i^{4n}$$

Complex Number :-

The number of the form  $a+ib$ .

where  $a$  &  $b$  are real numbers &  $i = \sqrt{-1}$  are known as complex number.

$$\begin{array}{ccc} & (a) + i(b) & \\ \swarrow & & \searrow \\ \text{real part} & & \text{imaginary part.} \end{array}$$

In complex number  $z = a + ib$ , the real numbers  $a$  &  $b$  are respectively known as real and imaginary part of  $z$  and we write.

$$\begin{aligned} \text{Re}(z) &= a \\ \& \text{ Im}(z) &= b \end{aligned}$$

Thus, the set  $\mathbb{C}$  of all complex number is given by

$$\mathbb{C} = \{ z : z = a + ib ; \text{ where } a, b \in \mathbb{R} \}$$

Purely Real and Purely Imaginary numbers :-

A complex number  $z$  is said to be.

- ① Purely real if  $\text{Im}(z) = 0$  (ex - 2, -3,  $\sqrt{3}$  etc)
- ② Purely imaginary if  $\text{Re}(z) = 0$  (ex -  $2i$ ,  $-7i$ ,  $\sqrt{3}i$ ,  $\frac{1}{7}i$  etc)

Conjugate of a complex number :-

The conjugate of a complex number  $z$  denoted by  $\bar{z}$  is the complex number obtained by changing the sign of - imaginary part of  $z$ .

ex -

$$\begin{aligned} \text{① } z &= 2 + 5i \\ \bar{z} &= (\overline{2 + 5i}) \\ &= 2 - 5i \end{aligned}$$

$$\begin{aligned} \text{③ } z &= 7i \\ \bar{z} &= (\overline{7i}) \\ &= -7i \end{aligned}$$

$$\begin{aligned} \text{② } z &= \sqrt{3} + 7i \\ \bar{z} &= (\overline{\sqrt{3} + 7i}) \\ &= \sqrt{3} - 7i \end{aligned}$$

$$\begin{aligned} \text{④ } z &= -9i \\ \bar{z} &= (\overline{-9i}) \\ &= 9i \end{aligned}$$

Modulus of a complex number :-

If  $z = x + iy$  be a complex number, the modulus of  $z$  written as  $|z|$  is a real number  $\sqrt{x^2 + y^2}$

$$\begin{aligned} \text{ex - ① } z &= 3 + 4i \\ |z| &= \sqrt{3^2 + 4^2} \\ &= 5 \end{aligned}$$

$$\begin{aligned} \text{② } z &= 6 + 2i \\ |z| &= \sqrt{6^2 + 2^2} \\ &= 2\sqrt{10} \end{aligned}$$

\* Also  $|z| = |\bar{z}|$



Equality of complex number :-

Two complex numbers  $z_1 = a_1 + ib_1$


&  $z_2 = a_2 + ib_2$  are said to be equal if their real part must be equal to  $z_1$ 's real part and imaginary part is must be equal to  $z_1$ 's - imaginary part that is  $a_1 = a_2$  and  $b_1 = b_2$ .  
That is

$$\text{Re}(z_1) = \text{Re}(z_2)$$

$$\& \text{Im}(z_1) = \text{Im}(z_2)$$

Geometrical representation of complex number :-

Complex numbers as Order pairs :-



We know that a complex number is of the form  $z = a + ib$  where  $a$  &  $b$  are real numbers thus, corresponding to each  $z = a + ib$  there is associated a unique order pair  $(a, b)$  of real numbers. So, we may represent

$$z = a + ib \text{ by } (a, b)$$

Thus, if  $z_1 = a, b$  and  $z_2 = c, d$  then we may define

$$z_1 + z_2 = (a, b) + (c, d)$$

$$= (a+c, b+d)$$

$$z_1 z_2 = (a+ib)(c+id)$$

$$= ac + iad + ibc + i^2 bd$$

$$= ac + (-1)bd + iad + ibc$$


$$= ac - bd + i(ad + bc)$$

clearly  $z_1 \Leftrightarrow z_2$

$$\Leftrightarrow (a, b) = (c, d) \quad \Leftrightarrow a = c, b = d$$

## Geometrical Representation :-

Let 'O' be the origin  $x'Ox$  &  $y'Oy$  be the co-ordinate axis. The real numbers are taken along x-axis and the imaginary numbers ~~are~~ taken along y axis. So, the



x axis is called the real axis and the y axis is called the imaginary axis. Then, any complex number  $z = a + ib$  may be represented by a unique point  $P(a, b)$  whose co-ordinates are  $(a, b)$  order pair. The ~~rep~~ - representation of a complex number as points in a plane forms an Argand diagram.

\* The plane on which complex numbers are represented is known as the complex plane or Argand's plane or Gaussian plane.

\* Let 'a' be a real number then we can write -  
$$a = a + i0$$
$$= (a, 0)$$

Let  $x'Ox$  and  $y'Oy$  be the co-ordinate axis.

Let  $z = a + ib$  be the complex number represented by point  $P = (a, b)$

Draw  $PM \perp OX$

then,

$OM = a$  and  $PM = b$  Join  $OP$

Let  $OP = r$  and  $\angle xOP = \theta$  then  $a = r \cos \theta$   
and  $b = r \sin \theta$ .

$$\begin{aligned}
 z &= a + ib \\
 &= r \cos \theta + i r \sin \theta \\
 &= r (\cos \theta + i \sin \theta)
 \end{aligned}$$

$$r = \sqrt{a^2 + b^2} = |z|$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{b}{r}}{\frac{a}{r}} = \frac{b}{a}$$

$$\theta = \tan^{-1} \left( \frac{b}{a} \right)$$

The form  $z = r (\cos \theta + i \sin \theta)$  is called the polar form or trigonometrical or standard form or modulus form or amplitude form of  $z$ .

Here,  $r = |z|$  and the angle  $\theta$  is known as amplitude or argument of  $z$ .

written as  $\text{amp}(z)$  or  $\text{arg}(z)$ .

The unique value of  $\theta$  such that  $-\pi < \theta \leq \pi$  for which  $a = r \cos \theta$  or  $b = r \sin \theta$  is known as the principle-value of amplitude.

The general value of amplitude is  $(2n\pi + \theta)$  where  $n$  is an integer and  $\theta$  is a principle value of.

Theorem :- 1

If  $z \in \mathbb{C}$  then

$$(1) \overline{\overline{z}} = z$$

$$(2) (z + \overline{z}) = 2 \text{Re}(z)$$

which is real

$$(3) (z - \overline{z}) = 2i \text{Im}(z)$$

which is an imaginary no.

$$(4) (z \overline{z}) = |z|^2 \text{ \& there for}$$

$z \overline{z}$  is real

$$(5) \text{Re}(z) \leq |z| \text{ \& } \text{Im}(z) \leq |z|$$

$$\text{Let } z = a + ib$$

$$\bar{z} = a - ib$$

$$\textcircled{1} \overline{(\bar{z})}$$

$$= \overline{(a - ib)}$$

$$= a + ib = z$$

$$\textcircled{2} (z + \bar{z})$$

$$= (a + ib) + (a - ib)$$

$$= a + \cancel{ib} + a - \cancel{ib}$$

$$= 2a = 2\operatorname{Re}(z)$$

$$\textcircled{3} (z - \bar{z})$$

$$= (a + ib) - (a - ib)$$

$$= \cancel{a} + ib - \cancel{a} + ib$$

$$= 2ib$$

$$\textcircled{4} (z\bar{z})$$

$$= (a + ib)(a - ib)$$

$$= (a)^2 - (ib)^2$$

$$= a^2 - i^2 b^2$$

$$= a^2 - (-1)^2 b^2$$

$$= a^2 + b^2 = |z|^2$$

$$\textcircled{5} \text{ Let } z = x + iy$$

$$|z| = \sqrt{x^2 + y^2}$$

$$x^2 \leq x^2 + y^2$$

$$\Rightarrow x \leq \sqrt{x^2 + y^2}$$

$$\text{i.e. } \operatorname{Re}(z) \leq |z|$$

$$y^2 \leq x^2 + y^2$$

$$\Rightarrow y \leq \sqrt{x^2 + y^2}$$

$$\text{i.e. } \operatorname{Im}(z) \leq |z|$$



## Theorem: 2

If  $z_1, z_2 \in \mathbb{C}$  then,

$$\textcircled{1} (z_1 + z_2) = \bar{z}_1 + \bar{z}_2$$

$$\textcircled{2} (z_1 - z_2) = \bar{z}_1 - \bar{z}_2$$

$$\textcircled{3} (\overline{z_1 z_2}) = \bar{z}_1 \bar{z}_2$$

$$\textcircled{4} |z_1 z_2| = |z_1| |z_2|$$

$$\textcircled{4} |z_1 z_2| = |z_1| \cdot |z_2|$$

$$\text{Let } z_1 = (a+ib) \text{ \& } z_2 = (c+id)$$

$$|z_1| = \sqrt{a^2 + b^2}$$

$$|z_2| = \sqrt{c^2 + d^2}$$

$$z_1 z_2 = (a+ib)(c+id)$$

$$= (ac - bd) + i(ad + bc)$$

Proof

$$\text{Let } z_1 = a+ib$$

$$\text{ \& } z_2 = c+id$$

$$\textcircled{1} z_1 + z_2 = (a+c) + i(b+d)$$

$$\overline{(z_1 + z_2)} = \overline{a+c + i(b+d)} = a+c - i(b+d)$$

$$= a+c - ib - id$$

$$= (a-ib) + (c-id)$$

$$= \bar{z}_1 + \bar{z}_2$$

[Proved]

$$\textcircled{2} z_1 - z_2 = (a-c) + i(b-d)$$

$$\overline{(z_1 - z_2)} = \overline{a-c + i(b-d)} = a-c - i(b-d)$$

$$= (a-ib) - (c-id)$$

$$= \bar{z}_1 - \bar{z}_2$$

[Proved]

$$\textcircled{3} \overline{(z_1 z_2)} = \bar{z}_1 \cdot \bar{z}_2$$

$$\text{Let } z_1 = (a+ib) \text{ \& } z_2 = (c+id)$$

$$z_1 z_2 = (a+ib)(c+id)$$

$$= (ac - bd) + i(ad + bc)$$

$$= (a-ib) \cdot (c-id)$$

$$\overline{z_1 z_2} = (ac - bd) - i(ad + bc)$$

$$= (a-ib) \cdot (c-id)$$

$$= \overline{(a+ib)} \cdot \overline{(c+id)}$$

$$= \bar{z}_1 \cdot \bar{z}_2$$

[Proved]

$$\begin{aligned} |z_1 z_2| &= \sqrt{\{(ac - bd)^2 + (ad + bc)^2\}} \\ &= \sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2} \\ &= |z_1| \cdot |z_2| \quad [\text{Proved}] \end{aligned}$$



Theorem :- 3

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

$$= (z_1 + z_2)(\bar{z}_1 + \bar{z}_2)$$

$$= z_1 \bar{z}_1 + z_1 \bar{z}_2 + z_2 \bar{z}_1 + z_2 \bar{z}_2$$

$$= |z_1|^2 + z_1 \bar{z}_2 + z_2 \bar{z}_1 + |z_2|^2$$

$$\boxed{z \bar{z} = |z|^2}$$

$$|z_1|^2 + |z_2|^2 + z_1 \bar{z}_2 + (z_2 \bar{z}_1)$$

$$= |z_1|^2 + |z_2|^2 + 2\operatorname{Re}(z_1 \bar{z}_2) \leq |z_1|^2 + |z_2|^2 + 2|z_1|^2 \left[ \because \operatorname{Re}(z) \leq |z| \right]$$

$$= |z_1|^2 + |z_2|^2 + 2|z_1||z_2|$$

$$= |z_1|^2 + |z_2|^2 + 2|z_1||z_2| \left[ \because |z| = |\bar{z}| \right]$$

$$= (|z_1| + |z_2|)^2$$

Reciprocal of a complex number :-

Let  $z = a + ib$  then the reciprocal of  $z = 1/z$

$$= \frac{1}{a + ib}$$

$$= \frac{1}{a + ib} \times \frac{a - ib}{a - ib}$$

$$= \frac{a - ib}{a^2 - i^2 b^2}$$

$$= \frac{a - ib}{a^2 + b^2}$$

$$= \left( \frac{a}{a^2 + b^2} \right) - i \left( \frac{b}{a^2 + b^2} \right)$$

Ex-1

$$z = a + ib$$

$$z = r(\cos \theta + i \sin \theta)$$

$$\Rightarrow 1 + i = r(\cos \theta + i \sin \theta)$$

$$\Rightarrow 1 + i = r \cos \theta + i r \sin \theta$$

$$r \cos \theta = 1$$

$$r \sin \theta = 1$$

$$r^2 \cos^2 \theta + r^2 \sin^2 \theta = 2$$

$$\Rightarrow r^2 = 2 \quad \Rightarrow r = \sqrt{2}$$

$$\cos \theta = 1/\sqrt{2}$$

$$\sin \theta = 1/\sqrt{2}$$

$$\Rightarrow \theta = \frac{\pi}{4}$$

$$1 + i = r(\cos \theta + i \sin \theta)$$

$$= \sqrt{2} \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$$

Hence the polar form of  $z = \sqrt{2} \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$

### Chapter - 3 Differential Equations

Solutions of a Differential Equations:-

The solution of a general ordinary differential equation of  $n$ th order.

$$F\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^n y}{dx^n}\right) = 0$$

$$\text{or } F(x, y, y', y'' \dots y^{(n)}) = 0 \dots \dots (1)$$

Definition:-

Let  $y = \phi(x)$ , define  $y$  as a real function of  $x$  on a real interval  $I$ . Then  $\phi$  is called an explicit sol<sup>n</sup> or simply a solution of the differential equation (1), and if we put  $y = \phi(x)$  and the given equation which is of the form,  $F(x, \phi(x), \phi'(x), \dots, \phi^{(n)}(x)) = 0$

Definition:-

A relation  $g(x, y) = 0$  is called implicit satisfied sol<sup>n</sup> of the differential equation by putting  $y = \phi(x)$  and ' $I$ ' such that  $\phi$  is an explicit sol<sup>n</sup>.

Any relation between the dependent variables not involving the derivatives which, when substituted in the differential eq<sup>n</sup> reduces it to an identity is called solution of the differential equation.

General Solution: (or complete primitive):-

The general solution of a differential equation is that in which the number of arbitrary constants is equal to the order of the differential equation and which satisfies the given differential equation.



# PRIMITIVE OR SOLUTION OF A DIFFERENTIAL EQUATION

A primitive or solution of a differential equation is a functional such that this relation and the derivatives - obtained from it satisfy the given differential equation.

For example,  $x = \cot y + c$  is the solution of the differential equation  $\frac{dy}{dx} + \sin^2 y = 0$

Now  $x = \cot y + c$  gives us

$$\frac{dy}{dx} = -\operatorname{cosec}^2 y$$

$$\text{or } \frac{dy}{dx} = -\sin^2 y$$

Substituting the value of  $\frac{dy}{dx}$  in L.H.S. of differential equation, we get

$$-\sin^2 y + \sin^2 y = 0$$

$$\text{L.H.S.} = \text{R.H.S.}$$

Thus the sol<sup>n</sup> of a differential eq<sup>n</sup> is a functional relation between  $x$  and  $y$  which is free from derivatives and this relation on substitution satisfy the differential eq<sup>n</sup>.

General sol<sup>n</sup> :-

It is that sol<sup>n</sup> which contains the number of arbitrary constants equal to the order of the differential eq<sup>n</sup>. It is also called complete primitive.

Thus in the above example the sol<sup>n</sup> contains one arbitrary constant and the eq<sup>n</sup> is of first order.

Particular sol<sup>n</sup> :-

A particular sol<sup>n</sup> of differential eq<sup>n</sup> is sol<sup>n</sup> obtained from the general sol<sup>n</sup> by giving particular values to the arbitrary constant. For example, putting  $c=1, 2$  etc, we have  $x = \cot y + 1$ ,  $x = \cot y + 2$  which are particular

Sol<sup>n</sup> of the eq<sup>n</sup>  $\frac{dy}{dx} + \sin^2 y = 0$

Note:- In exceptional cases a relation containing 'n' arbitrary constants may give rise to differential eq<sup>n</sup> of order less than n.

Formation of a differential eq<sup>n</sup> :-

We have seen that a general sol<sup>n</sup> to a differential eq<sup>n</sup> is a relation bet<sup>n</sup> the variables and it contains the number of arbitrary constants equal to the order of the eq<sup>n</sup>. As we have to study the differential eq<sup>n</sup> of the first order, let us have a functional relation

$$F(x, y, c) = 0 \quad \text{--- (1)}$$

Now we shall form that differential eq<sup>n</sup> whose sol<sup>n</sup> is

$$F(x, y, c) = 0$$

The required differential eq<sup>n</sup> will be obtained by eliminating 'c' from eq<sup>n</sup> (1) and another eq<sup>n</sup> obtained by differentiation (1) w.r.t x.

In other words we have to eliminate 'c' from the eq<sup>n</sup>s.

$$F(x, y, c) = 0$$

Eliminating 'c' from those two equations, we get the required differential eq<sup>n</sup>.



Example - 1

Q. Find the differential eq<sup>n</sup> of the family of curves  
 $y = e^x (A \cos x + B \sin x)$

Sol<sup>n</sup> given that the family of curves  $y = e^x (A \cos x + B \sin x)$   
 differentiate w.r.t  $x$ .

$$\frac{dy}{dx} = \frac{d(e^x)}{dx} (A \cos x + B \sin x) + e^x \left\{ \frac{d(A \cos x)}{dx} + \frac{d(B \sin x)}{dx} \right\}$$

$$= e^x (A \cos x + B \sin x) + e^x (-A \sin x + B \cos x)$$

$$\Rightarrow \frac{dy}{dx} = y + e^x (-A \sin x + B \cos x)$$

$$\Rightarrow \frac{dy}{dx} - y = e^x (-A \sin x + B \cos x)$$

again differentiate w.r.t  $x$

$$\frac{d^2y}{dx^2} - \frac{dy}{dx} = e^x (-A \sin x + B \cos x) + e^x \{-A \cos x + B \sin x\}$$

$$\Rightarrow \frac{d^2y}{dx^2} - \frac{dy}{dx} = e^x (-A \cos x + B \sin x) - y$$

$$\Rightarrow \frac{d^2y}{dx^2} - \frac{dy}{dx} + y = \frac{dy}{dx} - y$$

$$\Rightarrow \frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 2y = 0$$

which is the required ordinary differential eq<sup>n</sup>.

Sol<sup>n</sup> of the differential eq<sup>n</sup> of first order and first degree :-

Type 1 :

Equation of the type  $\frac{dy}{dx} = F(x)$

$$\frac{dy}{dx} = F(x)$$

$$\Rightarrow dy = F(x) \cdot dx$$

Integrating both side we get.

$$\Rightarrow \int dy = \int F(x) dx$$

$$\Rightarrow y = \phi(x) + c \text{ where } \phi(x) = \int F(x) dx$$

Type 2 :

Equation of the type  $\frac{dy}{dx} = F(y)$

$$\frac{dy}{dx} = F(y)$$

$$\Rightarrow \frac{dy}{F(y)} = dx$$

On Integrating

$$\Rightarrow \ln(F(y)) + c = x$$

$$\Rightarrow F(y) = e^x + c$$

Type 3 :

Equation with variable separable :-

If a given differential equation of a being expressed in the form  $F(x) \cdot dy + g(y) \cdot dx = 0$

$$\text{i.e. } \phi(x) dy + g(y) \cdot dx = 0$$

$$\phi(x) dy = -g(y) \cdot dx$$

$$\Rightarrow \frac{dy}{g(y)} = \frac{-dx}{\phi(x)}$$

on integrating

$$\Rightarrow \ln(g(y)) = -\ln(F(x)) + \ln c$$

$$\Rightarrow \ln(g(y)) + \ln(F(x)) = \ln c$$

$$\Rightarrow \ln(g(y) F(x)) = \ln c$$

where  $g(y)$  &  $F(x)$  are respectively functions of  $y$  &  $x$  is called a variable and Separable

The sol<sup>n</sup> of such an eq<sup>n</sup> is obtain by Integrating each term separately.

Type - 4 :

Solving of differential eq<sup>n</sup> of second order of

$$\frac{d^2 y}{dx^2} = \psi(x)$$

$$\Rightarrow \frac{d}{dx} \left( \frac{dy}{dx} \right) = \psi(x)$$

on integration, both side

$$\Rightarrow \frac{dy}{dx} = \int \psi(x) \cdot dx + c_1$$

$$\text{let } \int \psi(x) \cdot dx = \phi(x)$$

$$\Rightarrow \frac{dy}{dx} = \phi(x) + c_1$$

now multiplying both side with differential  $dx$  and the the integrating both side.

$$\Rightarrow \int dy = \int (\phi(x) + c_1) \cdot dx$$

$$\Rightarrow y = \int \phi(x) dx + c_2 + c_1 x$$

$$\Rightarrow y = \int \phi(x) dx + c_1 x + c_2$$

$$= \psi(x) + c_1 x + c_2$$

where  $\psi(x) = \int \phi(x) dx$ .  
 $c_2$  is the constant integration which is independent  
 $c_1$ .

$y = \psi(x) + c_1x + c_2$  is the required differential eqn.

Type-5:-

Equation reducible to variable separable

The eqn of the form  $\frac{dy}{dx} = f(ax+by+c)$  can be  
reduce to variable separable by the substitution  
 $ax+by+c = z$

Type 6:-

Homogeneous function.

A function  $f(x, y)$  in  $x$  and  $y$  is called a  
homogeneous function of degree  $n$ , if the degree of  
each term is  $n$ .

Example - 1 :-

$f(x, y) = x^2 + y^2 - xy$  is a homogeneous function of  
degree 2.

Example - 2 :-

$f(x, y) = x^3 + 5x^2y + x^2y$  is a homogeneous fun<sup>n</sup> of  
degree 3.

Degree of Homogeneous differential eqn:-

A differential eqn of the form  $\frac{dy}{dx} = \frac{f(x, y)}{g(x, y)}$

where  $f(x, y)$  as well as  $g(x, y)$  is a homogeneous  
fun<sup>n</sup> of same degree in  $x$  and  $y$  is called a  
homogeneous differential eqn.



Ex

$$\textcircled{1} \frac{dy}{dx} = \frac{x^2 + y^2}{2xy}$$

$\textcircled{2} \frac{dy}{dx} = x \cos(y/x)$  is a homogeneous differential eqn as  $x \cos(y/x)$  being a fun<sup>n</sup> of  $y/x$  is a homogeneous fun<sup>n</sup>.

Method of solving a homogeneous eqn :-

Let  $\frac{dy}{dx} = f(x, y) g(x, y)$  is a homogeneous differential eqn.

Put  $y = vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

Substitute this values in the given eqn we get  
~~v~~  $v + x \frac{dv}{dx} = f(v)$

$$\Rightarrow x \frac{dv}{dx} = f(v) - v$$

$$\Rightarrow \frac{dv}{f(v) - v} = \frac{dx}{x}$$

On integration both side

$$\Rightarrow \int \frac{dv}{f(v) - v} = \int \frac{dx}{x}$$

$$\Rightarrow \ln + C$$

Now replace  $v$  by  $y/x$  to obtain the require sol<sup>n</sup>.



Type 7 :

Equation reducible to homogeneous function.  
Equation of the type.

$$\frac{dy}{dx} = \frac{ax + by + c}{Ax + By + c}$$

$$x = x + h$$

$$y = y + k$$

where  $h$  &  $k$  are constant.

$$\frac{dy}{dx} = \cancel{d(x+y)} \cdot \frac{d(y+k)}{dx} = \frac{d(y+k)}{dx} \quad \frac{dx}{dx} \uparrow = \frac{dy}{dx}$$

$$\frac{ax + by + c}{Ax + By + c}$$

$$= \frac{a(x+h) + b(y+k) + c}{A(x+h) + B(y+k) + c}$$

$$= \frac{ax + by + (ah + bk + c)}{Ax + By + (Ah + Bk + c)}$$

$$\begin{cases} ah + bk + c = 0 \\ Ah + Bk + c = 0 \end{cases}$$

$$\Rightarrow Ah + Abk + Ac = 0$$

$$\begin{matrix} (-) & & (-) & & (-) \\ Ah + Abk + Ac = 0 \end{matrix}$$

$$Abk - abk = ac - ca$$

$$\Rightarrow k = \frac{ac - ca}{Ab - aB}$$

Exact Equation :-

The differential eq<sup>n</sup>  $M(x, y) dx + N(x, y) dy = 0$  is exact if and only  $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$  where,  $\frac{\partial M}{\partial y}$  denotes the differential co-efficient of  $M$  with respect to  $y$  keeping  $x$  constant.

Rule-1

Rule-2

~~Integ~~ Integrate w.r.t  $y$ . only those terms of  $N$  which do not contain  $x$ .

Rule-3

Result of ① + Result of ② = constant

Integrating Factor :-

An integrating factor is a function when multiplied by it the left hand side of eq<sup>n</sup>  $M(x, y) dx + N(x, y) dy = 0$  becomes the exact equation.

$M(x, y) dx + N(x, y) dy = 0$  — ① is not exact or total differential.  
It is easy to choose fun<sup>n</sup>  $\mu(x, y)$  such the after multiplying by it the left side of eq<sup>n</sup> ① becomes the exact differential  $du = \mu(x, y) M(x, y) dx + \mu(x, y) N(x, y) dy$

Note:-

① The number of integrating factor is infinite.

② If  $Mx + Ny \neq 0$  and eq<sup>n</sup> ① is homogenous then

$\frac{1}{Mx + Ny}$  is the integrating factor of the ~~integrating~~ integration.

$$\textcircled{3} \quad Mx + Ny = 0$$

$$\Rightarrow Mx = -Ny$$

$$\Rightarrow \frac{M}{N} = \frac{-y}{x}$$

$\textcircled{4}$  If  $Mx - Ny \neq 0$  and the eqn —  $\textcircled{1}$  has the form  
 $M(x, y) dx + N(x, y) dy = 0$   
 $\frac{1}{Mx - Ny}$  is the integrating factor.

Linear differential eqn :-

A differential eqn in which the dependent variables and the both occur in the first degree only and are not multiplied together, is called a linear differential eqn.

$$\frac{dy}{dx} + 5y = x^2 \text{ is a linear eqn of order 1.}$$

$$x \frac{d^2x}{dy^2} + 5 \frac{dy}{dx} = 8 \text{ is a linear eqn of order degree 2.}$$

$$x \left( \frac{dy}{dx} \right)^2 - y^2 \frac{dy}{dx} + 3 = 0 \text{ is a Non-linear differential eqn of order 2 degree 2.}$$

Note :-

Every linear differential eqn is of degree 1 but every differential eqn of degree 1 is need not linear.

Type 1 :

$\frac{dy}{dx} + py = Q$ , where  $p$  is a constant and  $Q$  may be a constant or a fun<sup>n</sup> of  $x$  only.

$$\text{To solve } \frac{dy}{dx} + py = Q,$$

First we find  $\int p dx$ , which is known as Integrating factor, written as I.F.

Multiplying both sides of the given eqn by  $\int p dx$  we get

$$\int p dx \cdot \frac{dy}{dx} + p y \int p dx = Q \int p dx$$

or,

$$\int p dx \frac{dy}{dx} + p y \int p dx = Q \int p dx dx$$

or,

$$\int \frac{d}{dx} (y \int p dx) = Q \int p dx dx$$

Integrating we get

$$y \cdot \int p dx = \int Q \int p dx dx + c$$

which is the required sol<sup>n</sup> of the given diff<sup>n</sup> eqn.  
working Rule for solving  $\frac{dy}{dx} + py = Q$

(i) Find I.F =  $\int p dx$

(ii) The sol<sup>n</sup> is  $y \times (I.F)$

$$= \int \{Q \times (I.F)\} dx + c$$

Type 2:

Differential eqns linear in  $x$  and  $\frac{dx}{dy}$

These eqns are of the form  $\frac{dx}{dy} + px = Q$

where  $p$  and  $Q$  are fcn's of  $y$  only or constants



The sol<sup>n</sup> is given by Integrating factor = I.F. =  $\int P dy$   
Next to find it's sol<sup>n</sup>

$$\propto \int P dy = \int Q \cdot \int P dy \cdot dy + C$$

$$\propto (I.F) = \int Q \cdot (I.F) dy + C$$

Type 3:

Equations reducible to the linear form:

$$(a) \frac{dy}{dx} + Py = Qy^n$$

where  $P$  and  $Q$  are constants or fun<sup>n</sup>s of  $x$  alone and  $n$  is a constant other than zero or unity can be reduced to the linear form by  $y^n$  and

Substituting  $\frac{1}{y^{n-1}} = z$

on dividing the given eq<sup>n</sup> by  $y^n$ , we get

$$\frac{1}{y^n} \left( \frac{dy}{dx} \right) + \frac{1}{y^{n-1}} P = Q$$

Put  $\frac{1}{y^{n-1}} = z$ , then  $\frac{-n+1}{y^n} \frac{dy}{dx} = \frac{dz}{dx}$  and

(2) becomes  $\frac{1}{(-n+1)} \frac{dz}{dx} + Pz = Q$ , which is linear eq<sup>s</sup>:

(b) Similarly  $F'(y) \frac{dy}{dx} + PF(y) = Q$ , can be reduced to the linear form by the substitution  $F(y) = z$ .



Example 1:

$$\text{Solve } \frac{dy}{dx} + xy = xy^3$$

Dividing by  $y^3$ , the eq<sup>n</sup> becomes

$$\Rightarrow \frac{dy}{dx} + \frac{3}{x} y = x$$

$$\text{where } p = \frac{3}{x} \text{ \& } Q = x$$

$$\begin{aligned} \text{I.F} &= e^{\int p \cdot dx} = e^{\int \frac{3}{x} \cdot dx} = e^{3 \ln x + c} \\ &= e^{\ln x^3} = x^3 + c \end{aligned}$$

So the required sol<sup>n</sup> is

$$\text{I.F } xy = \int (Q \times \text{I.F}) dx + c$$

$$\Rightarrow x^3 xy = \int (x \times x^3) dx + c$$

$$\Rightarrow x^3 y = \frac{x^5}{5} + c$$

Example 2:

$$\text{Solve } \frac{dy}{dx} + (\sec x) y = \tan x$$

clearly from the above differential eq<sup>n</sup>

we get  $p = \sec x$  \&  $Q = \tan x$

$$\begin{aligned} \text{Then, I.F} &= e^{\int p dx} = e^{\int \sec x \cdot dx} = e^{\ln |\sec x + \tan x|} + c \\ &= (\sec x + \tan x) + c \end{aligned}$$

So, the required sol<sup>n</sup> is

$$\text{I.F } xy = \int Q + \text{I.F } dx + c$$

$$y(\sec x + \tan x) = \int \tan x (\sec x + \tan x) dx + C$$

$$y(\sec x + \tan x) = \int \sec x \tan x dx + \int \sec^2 x dx + C$$

$$\text{or } y(\sec x + \tan x) = \sec x + \tan x - x + C.$$

Date - 15.12.2022

Ex-1

$$(x + 2y^3) \frac{dy}{dx} = y$$

$$\Rightarrow x + 2y^3 = y \cdot \frac{dx}{dy}$$

$$\Rightarrow y \cdot \frac{dx}{dy} - x = 2y^3$$

$$\Rightarrow \frac{dx}{dy} - \frac{x}{y} = 2y^2$$

$$P = -1/y \text{ \& } Q = 2y^2$$

$$\text{I.F} = e^{\int P \cdot dy}$$

$$= e^{\int -1/y dy}$$

$$= e^{\int 1/y \cdot dy}$$

$$= e^{-\ln y}$$

$$= e^{\ln y^{-1}}$$

$$= y^{-1}$$

$$= 1/y$$

Thus, the soln is

$$\text{I.F} \cdot x = \int (Q \times \text{I.F}) dy + C$$

$$\begin{aligned}
 \Rightarrow \frac{x}{y} &= \int (2y^2 \times \frac{1}{y}) dy + c \\
 &= 2 \int y \cdot dy + c \\
 &= y^2 + c \\
 \Rightarrow x &= y(y^2 + c)
 \end{aligned}$$

Ex-2

$$(1+y^2) dx = (\tan^{-1} y - x) dy$$

$$\Rightarrow (1+y^2) \frac{dx}{dy} = \tan^{-1} y - x$$

$$\Rightarrow (1+y^2) \frac{dx}{dy} + x = \tan^{-1} y$$

$$\Rightarrow \frac{dx}{dy} + \frac{x}{1+y^2} = \frac{\tan^{-1} y}{1+y^2}$$

$$P = \frac{1}{1+y^2}$$

$$Q = \frac{\tan^{-1} y}{1+y^2}$$

$$\text{I.F.} \cdot \int P \cdot dy = \int \frac{1}{1+y^2} dy = e^{\tan^{-1} y}$$

Thus the sol<sup>n</sup> is

$$\text{I.F.} \times x = \int (Q \times \text{I.F.}) dy + c$$

$$x \cdot e^{\tan^{-1} y} = \int \left( \frac{\tan^{-1} y}{1+y^2} \times e^{\tan^{-1} y} \right) dy + c$$

$$\left[ \begin{array}{l} \text{Put } z = \tan^{-1} y \\ \frac{dy}{1+y^2} = dz \end{array} \right]$$

$$\begin{aligned}
 \Rightarrow x \cdot e^z &= \int z \cdot e^z \cdot dz + c \\
 &= \int z e^z - e^z + c \\
 &= x \cdot e^z = e^z (z-1) + c
 \end{aligned}$$

① For any  $z_1, z_2 \in \mathbb{C}$ , prove that

$$\textcircled{1} |z_1 + z_2| \leq |z_1| + |z_2|$$

$$\textcircled{2} |z_1 - z_2| \leq |z_1| + |z_2|$$

$$\text{A. } \textcircled{1} |z_1 + z_2|^2 = (z_1 + z_2) (\overline{z_1 + z_2})$$

$$= (z_1 + z_2) (\overline{z_1} + \overline{z_2})$$

$$= z_1 \overline{z_1} + z_2 \overline{z_2} + z_1 \overline{z_2} + z_2 \overline{z_1}$$

$$= |z_1|^2 + |z_2|^2 + z_1 \overline{z_2} + \overline{z_1} z_2$$

$$= |z_1|^2 + |z_2|^2 + 2\operatorname{Re}(z_1 \overline{z_2}) \leq |z_1|^2 + |z_2|^2 + 2|z_1| |z_2|$$

$$= |z_1|^2 + |z_2|^2 + 2|z_1| |z_2|$$

$$= (|z_1| + |z_2|)^2$$

$$\therefore |z_1 + z_2| \leq |z_1| + |z_2|$$

$$\textcircled{2} |z_1 - z_2| = |z_1 + (-z_2)| \leq |z_1| + |-z_2|$$

$$= |z_1| + |z_2|$$

$$\therefore |z_1 - z_2| \leq |z_1| + |z_2|$$

$$\textcircled{2} |z_1 z_2| = |z_1| |z_2| \text{ \& } \arg |z_1 z_2| \neq \arg(z_1) + \arg(z_2)$$

$$\text{+ let } z_1 = r_1 (\cos \theta_1 + i \sin \theta_1)$$

$$z_2 = r_2 (\cos \theta_2 + i \sin \theta_2)$$

Then,

$$z_1 z_2 = r_1 r_2 (\cos \theta_1 + i \sin \theta_1) (\cos \theta_2 + i \sin \theta_2)$$

$$= r_1 r_2 [(\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2) + (i \sin \theta_1 \cos \theta_2 + \cos \theta_1 i \sin \theta_2)]$$



$$= r_1 r_2 \left[ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \right]$$

$$|z_1 z_2| = r_1 r_2 = |z_1| |z_2|$$

$$\text{and } \arg(z_1 z_2) = \theta_1 + \theta_2 = \arg(z_1)$$

$$\text{Ex (ii)} \quad \frac{z_1}{z_2} = \frac{r_1 (\cos \theta_1 + i \sin \theta_1)}{r_2 (\cos \theta_2 + i \sin \theta_2)} \times \frac{(\cos \theta_2 - i \sin \theta_2)}{(\cos \theta_2 - i \sin \theta_2)}$$

$$= \frac{r_1}{r_2} \frac{(\cos \theta_1 + i \sin \theta_1) [\cos(-\theta_2) + i \sin(-\theta_2)]}{[\cos^2 \theta_2 + \sin^2 \theta_2]}$$

$$= \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]$$

$$\therefore \left| \frac{z_1}{z_2} \right| = \frac{r_1}{r_2} = \frac{|z_1|}{|z_2|}$$

$$\text{and } \arg \left| \frac{z_1}{z_2} \right| = (\theta_1 - \theta_2) = [\arg(z_1) - \arg(z_2)]$$

$$\text{③ If } z_1, z_2 \in \mathbb{C}, \text{ then } \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$$

4. Let,

$$z_1 = a + ib$$

$$z_2 = c + id$$

$$\frac{z_1}{z_2} = \frac{a+ib}{c+id} = \frac{a+ib}{c+id} \times \frac{c-id}{c-id}$$

$$= \frac{(a+ib)(c-id)}{c^2 + d^2}$$

$$= \frac{(ac + bd) + i(bc - ad)}{c^2 + d^2}$$

$$= \frac{ac + bd}{c^2 + d^2} + i \frac{bc - ad}{c^2 + d^2}$$

$$\left| \frac{z_1}{z_2} \right| = \sqrt{\left( \frac{ac+bd}{c^2+d^2} \right)^2 + \left( \frac{bc-ad}{c^2+d^2} \right)^2}$$

$$= \sqrt{\frac{a^2c^2 + b^2d^2 + b^2c^2 + a^2d^2}{(c^2+d^2)^2}}$$

$$= \sqrt{\frac{(a^2+b^2)(c^2+d^2)}{(c^2+d^2)^2}}$$

$$= \sqrt{\frac{a^2+b^2}{c^2+d^2}}$$

$$= \frac{\sqrt{a^2+b^2}}{\sqrt{c^2+d^2}}$$

$$= \frac{|z_1|}{|z_2|}$$

Hence  $\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$

$$\textcircled{5} \quad \frac{(x+i)^2}{x-i} - \frac{(x-\sqrt{-1})^2}{x+i}$$

$$= \frac{(x+i)^2}{x-i} - \frac{(x-i)^2}{x+i}$$

$$= \frac{(x+i)^3 - x(x-i)^3}{(x-i)(x+i)}$$

$$= \frac{\{(x+i)(x-i)(x+i)^2 + (x-i)^2 + (x+i)(x-i)\}}{x^2 - i^2}$$

$$= \frac{2i \{ x^2 + i^2 + 2ix + x^2 + i - 2xi + x^2 - i^2 \}}{x^2 + 1}$$

$$= \frac{2i(3x^2 + i^2)}{x^2 + 1}$$

$$= \frac{6x^2 i + 2i^3}{x^2 + 1}$$

$$= \frac{(6x^2 - 2)i}{x^2 + 1}$$

$$\textcircled{6} (1-\omega)(1-\omega^2)(1-\omega^4)(1-\omega^5) = 9$$

† LHS

$$\begin{aligned} & (1-\omega)(1-\omega^2)(1-\omega^4)(1-\omega^5) \\ &= (1-\omega)(1-\omega^2)(1-\omega^3 \cdot \omega)(1-\omega^3 \cdot \omega^2) \\ &= (1-\omega)(1-\omega^2)(1-\omega)(1-\omega^2) \\ &= (1-\omega^2)(1-\omega^2)^2 \\ &= \{ (1-\omega)(1-\omega^2) \}^2 \\ &= (1-\omega-\omega^2+\omega^3)^2 \\ &= (1-\omega-\omega^2+1)^2 \\ &= (2-\omega-\omega^2)^2 \\ &= (2+1)^2 = 3^2 = 9 \quad \underline{\underline{RHS}} \end{aligned}$$

⑩ Find the rank of the given matrix.

$$\begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{bmatrix}$$

A. Let  $A = \begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{bmatrix}$

$$\sim \begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 0 & 0 \\ 3 & 1 & -3 & -1 \\ 1 & 1 & -3 & -1 \end{bmatrix} \quad [C_3 \rightarrow C_3 - C_1, C_4 \rightarrow C_4 - C_1]$$

$$\sim \begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 0 & 0 \\ 3 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} R_3 \rightarrow R_3 - R_1 \\ R_4 \rightarrow R_4 - R_1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} R_3 \rightarrow R_3 - 3R_2 \\ R_4 \rightarrow R_4 - R_2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} C_3 \rightarrow C_3 + 3C_2 \\ C_4 \rightarrow C_4 + C_2 \end{bmatrix}$$

Hence rank of the matrix is 2.



⑪ Find the rank of the following matrices

$$\begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 4 \end{bmatrix} \quad \begin{bmatrix} R_2 \rightarrow R_2 + 2R_1 \\ R_3 \rightarrow R_3 + R_1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 0 \end{bmatrix} \quad \left[ R_3 \rightarrow R_3 - \frac{1}{2}R_2 \right]$$

Hence rank of matrix is 2.

⑫ Solve  $x + 2y - z = 3$   
 $3x - y + 2z = 1$   
 $2x - 2y + 3z = 2$

$$\sim \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \\ 2 & -2 & 3 & 2 \end{bmatrix}$$

$$R_2 = R_2 - 3R_1$$

$$\begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & -7 & 5 & -8 \\ 2 & -2 & 3 & 2 \end{bmatrix}$$

$$R_3 = R_3 - 2R_1$$

$$\sim \begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & -7 & 5 & -8 \\ 0 & -6 & 5 & -4 \end{bmatrix}$$

$$(C_2 \leftrightarrow C_3)$$

$$\sim \begin{bmatrix} 1 & -2 & 2 & 3 \\ 0 & 5 & -7 & -8 \\ 0 & 5 & -6 & -4 \end{bmatrix}$$

$$R_3 \rightarrow R_3 - R_2$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 3 \\ 0 & 5 & -7 & -8 \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

$\rho(A) = 3$  Obviously rank of  $P(K)$  matrix is also 3  
 $\rho(A) = \rho(K) = \text{eqns}$  are ~~consisten~~ consistent.

Next to find its ~~exp~~ sol<sup>n</sup>.

$$x - y + 2z = 3, 5y - 7z = -8, z = 4$$

By backward substitution we get  $z = 4$

$$5y - 7z = -8, 5y - 28 = -8, 5y = 20, y = 4$$

$$x - 4 + 8 = 3, x + 4 = 3, x = -1$$

or  $x = -1, y = 4, z = 4$  be the required solutions.

# Rules For Finding the complementary Function :-

To solve the equation

$$\frac{d^n y}{dx^n} + K_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + K_n y = 0 \dots \dots \textcircled{i}$$

where  $K_s$  are constants. The eqn (i) in symbolic form is

$$(D^n + K_1 D^{n-1} + K_2 D^{n-2} + \dots + K_n) y = 0$$

It's symbolic co-efficients are equated to zero i.e.

$$D^n + K_1 D^{n-1} + K_2 D^{n-2} + \dots + K_n = 0 \dots \dots \textcircled{ii}$$

is called auxiliary eqn (A.E) or characteristic eqn.

Let  $y = e^{mx}$  be the soln of eqn (ii), then substituting

$$y = e^{mx}, Dy = m e^{mx}, D^2 y = m^2 e^{mx}$$

$D^{n-1} = m^{n-1} e^{mx}$ ,  $D^n y = m^n e^{mx}$  in eqn (ii), we get

$$(m^n + K_1 m^{n-1} + K_2 m^{n-2} + \dots + K_{n-1} m + K_n) e^{mx} = 0$$

Since  $y = e^{mx}$  is a soln of (ii)

$$m^n + K_1 m^{n-1} + \dots + K_n = 0$$

i.e if 'm' is a root of the eqn, then

$$D^n + K_1 D^{n-1} + \dots + K_{n-1} D + K_n = 0$$

Let  $D = m_1, m_2, \dots, m_n$  be the roots of the auxiliary eqn.

Case 1: The roots of the A.E are all real and different.

Since the roots  $m_1, m_2, \dots, m_n$  of the A.E are all real and different then eqn (ii) may be written as

$$(D - m_1)(D - m_2) \dots (D - m_n) y = 0 \dots \dots \textcircled{iii}$$

Since eqn (iii) will be satisfied by the soln of the eqns.

$$(D - m_1) y = 0, (D - m_2) y = 0 \dots \dots (D - m_n) y = 0 \dots \dots \textcircled{iii}$$

Let us consider the eqn

$$(D - m_1) y = 0 \Rightarrow \frac{dy}{dx} - m_1 y = 0 \dots \dots \textcircled{iv}$$

This is Leibnitz is linear eqn, I.F =  $\int m_1 dx = e^{-m_1 x}$

Thus the soln of the eqn (iv) is  $y \cdot e^{-m_1 x} = \int 0 \cdot e^{-m_1 x} dx = \int 0 dx = C_1$   
 $y = C_1 e^{m_1 x}$

Similarly considering other factors  $y = C_2 e^{m_2 x}$ ,  $y = C_3 e^{m_3 x}$ , ...,  $y = C_n e^{m_n x}$   
Hence the complete soln is  $y = C_1 e^{m_1 x} + C_2 e^{m_2 x} + \dots + C_n e^{m_n x}$  ... (v)

Case 2: The two of the roots of the auxiliary eqn are equal:

$$\text{Let } m_1 = m_2$$

Then the soln obtained in (v) is not a general soln because  
In this case the result (v) becomes

$$y = (C_1 + C_2 x) e^{m_1 x} + C_3 e^{m_3 x} + \dots + C_n e^{m_n x}$$

Thus there are  $(n-1)$  arbitrary constants and not 'n'  
To find the general soln let us consider  $(D - m_1)^2 y = 0$

$$\text{Let } (D - m_1) y = V$$

Then the eqn reduces to

$$(D - m_1) V = 0 \text{ i.e. } \frac{dV}{dx} - m_1 V = 0$$

whose soln is  $V = C_1 e^{m_1 x}$

Substituting this value  $\frac{dy}{dx} - m_1 y = C_1 e^{m_1 x}$  of  $V$  as

$(D - m_1) y$ , we get

The eqn is in Leibnitz's form, I.F. =  $e^{-\int m_1 dx} = e^{-m_1 x}$

Next to find it's soln

$$y (\text{I.F.}) = \int (\text{I.F.}) Q \cdot dx \Rightarrow y \cdot e^{-m_1 x} = \int e^{-m_1 x} \cdot C_1 e^{m_1 x} dx$$

$$y \cdot e^{-m_1 x} = C_1 \int dx = C_1 x + C_2$$

Hence it's complete soln is  $y = (C_1 x + C_2) e^{m_1 x} + C_3 e^{m_2 x} + \dots + C_n e^{m_n x}$

However in this case three roots of the A.E are equal.

$$\text{Say } m_1 = m_2 = m_3$$

then proceeding as above the soln becomes

$$y = (C_1 x^2 + C_2 x + C_3) e^{m_1 x} + \dots + C_n e^{m_n x}$$



Case 3: Two of the roots of the ~~aux~~ auxiliary eqn are complex.

Let the two roots of A.E be  $m_1 = \alpha + i\beta$ ,  $m_2 = \alpha - i\beta$ , then the complete soln is

$$y = c_1 e^{(\alpha + i\beta)x} + c_2 e^{(\alpha - i\beta)x} + c_3 e^{m_3 x} + \dots + c_n e^{m_n x}$$

$$\begin{aligned} y &= e^{\alpha x} (c_1 e^{i\beta x} + c_2 e^{-i\beta x}) + c_3 e^{m_3 x} + \dots + c_n e^{m_n x} \\ &= e^{\alpha x} [c_1 (\cos \beta x + i \sin \beta x) + c_2 (\cos \beta x - i \sin \beta x)] + c_3 e^{m_3 x} + \dots + c_n e^{m_n x} \\ &= e^{\alpha x} [(c_1 + c_2) \cos \beta x + i(c_1 - c_2) \sin \beta x] + c_3 e^{m_3 x} + \dots + c_n e^{m_n x} \end{aligned}$$

By Euler's theorem  $e^{i\theta} = \cos \theta + i \sin \theta$

$$= e^{\alpha x} (A \cos \beta x + B \sin \beta x) + c_3 e^{m_3 x} + \dots + c_n e^{m_n x}$$

which is the required soln.

Case 4:

Let  $m_1 = m_2 = \alpha + i\beta$ ,  $m_3 = m_4 = \alpha - i\beta$

The complete soln is

$$y = e^{\alpha x} [(c_1 x + c_2) \cos \beta x + (c_3 x + c_4) \sin \beta x] + c_5 e^{m_5 x} + \dots + c_n e^{m_n x}$$

working Rule to solve the eqn :-

$$\frac{d^n y}{dx^n} + K_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + K_n y = X$$

of which the eqn is in symbolic form

$$D^n y + K_1 D^{n-1} y + \dots + K_n y = X$$

$$(D^n + K_1 D^{n-1} + \dots + K_n) y = X$$

$$f(D) y = X$$

$$\text{where } f(D) = (D^n + K_1 D^{n-1} + \dots + K_n)$$

Step 1:

To find the complementary fun<sup>n</sup>  
 (i) write the A.E  $F(D)y = 0$  and solve it. Let it's root be  $D = m_1, m_2, \dots, m_n$ .  
 write the complete sol<sup>n</sup>

Roots of A.E.

- (1) All the roots  $m_1, m_2, m_3, \dots, m_n$  are real & different.
- (2) All the roots  $m_1, m_2, m_3, \dots, m_n$  are real &  $m_1 = m_2$ .
- (3) All the roots  $m_1, m_2, m_3, \dots$  are real &  $m_1 = m_2 = m_3$ .
- (4) Two pairs of the roots are complex i.e.  $m_1 = \alpha + i\beta, m_2 = \alpha - i\beta$  & all other roots are real & different.

Complete solution:

- (i)  $y = c_1 e^{m_1 x} + c_2 e^{m_2 x} + c_3 e^{m_3 x} + \dots + c_n e^{m_n x}$
- (ii)  $y = (c_1 x + c_2) e^{m_1 x} + c_3 e^{m_3 x} + \dots + c_n e^{m_n x}$
- (iii)  $y = (c_1 x + c_2 x^2 + c_3) e^{m_1 x} + c_4 e^{m_4 x} + \dots + c_n e^{m_n x}$
- (iv)  $y = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x) + c_3 e^{m_3 x} + \dots + c_n e^{m_n x}$
- (v)  $y = e^{\alpha x} [(c_1 x + c_2) \cos \beta x + (c_3 x + c_4) \sin \beta x] + c_5 e^{m_5 x} + \dots + c_n e^{m_n x}$

Observe the following illustrative table for writing the complementary fun<sup>n</sup> based on the roots of the auxiliary eq<sup>n</sup>.

Roots of the A.E.

1. 2, 3
2. 1, -1, 2, -2
3. 3, 3
4. 0, -2, -2, -2
5.  $\pm 2i$
6.  $3 \pm 2i$
7. 1, 1,  $1 \pm i$
8.  $1 \pm 2i, 1 \pm 2i$

complementary fun<sup>n</sup> (y<sub>c</sub>)

- $c_1 e^{2x} + c_2 e^{3x}$
- $c_1 e^x + c_2 e^{-x} + c_3 e^{2x} + c_4 e^{-2x}$
- $(c_1 + c_2 x) e^{3x}$
- $c_1 e^{0x} + (c_2 + c_3 x + c_4 x^2) e^{-2x}$
- $c_1 \cos 2x + c_2 \sin 2x$
- $e^{3x} (c_1 \cos 2x + c_2 \sin 2x)$
- $(c_1 + c_2 x) e^x + e^x (c_3 \cos x + c_4 \sin x)$
- $e^x \{ (c_1 + c_2 x) \cos 2x + (c_3 + c_4 x) \sin 2x \}$

working procedure for problems to solve a homogeneous D.E with constant coefficients.

- The given D.E is put in the form  $F(D)y = 0$
- Form the A.E  $F(m) = 0$  and solve the same (Here we need to adopt various techniques to solve the eqn including the synthetic division method.)
- Based on the nature of the roots of the A.E we write the c.f which itself is the general soln of the D.E taking into account the dependent and the independent variables involved in the D.E.

Roots of the A.E.

1. 2, 3
2. 1, -1, 2, -2
3. 3, 3
4. 0, -2, -2, -2
5.  $\pm 2i$
6.  $3 \pm 2i$
7. 1, 1,  $1 \pm i$
8.  $1 \pm 2i, 1 \pm 2i$

complementary fun<sup>n</sup> ( $y_c$ )

$$\begin{aligned}
 & c_1 e^{2x} + c_2 e^{3x} \\
 & c_1 e^x + c_2 e^{-x} + c_3 e^{2x} + c_4 e^{-2x} \\
 & (c_1 + c_2 x) e^{3x} \\
 & c_1 e^{0x} + (c_2 + c_3 x + c_4 x^2) e^{-2x} \\
 & c_1 \cos 2x + c_2 \sin 2x \\
 & e^{3x} (c_1 \cos 2x + c_2 \sin 2x) \\
 & (c_1 + c_2 x) e^x + e^x (c_3 \cos x + c_4 \sin x) \\
 & e^x \{ c_1 + c_2 x \} \cos 2x + (c_3 + c_4 x) \sin 2x
 \end{aligned}$$



Solution of Differential Equation  $F(D)y = x$  or P.I.

We have already discuss from the previous chapter, that the sol of the equation  $F(D)y = x$ , consists of two parts, namely complementary function and particular integral. The complementary function for this equation is same as the complete sol<sup>n</sup> of  $F(D)y = 0$ . The methods to find complementary functions have already been discussed. we shall discuss the methods of finding the particular integrals.

**Particular Integral :-**

The particular integral of the differential equation

$$F(D)y = x \text{ is defined as } y = \frac{1}{F(D)} x \dots \dots \textcircled{1}$$

where in general 'x' is a function of  $x$ , it may or of course be constant. The symbol  $\frac{1}{F(D)} x$ , is defined as a function of  $x$ , which when separated upon by  $F(D)$ , gives  $x$ , in -  
Symbolically  $F(D) \left[ \frac{1}{F(D)} x \right] = x$ .

Thus the function  $\frac{1}{F(D)} x$  satisfies the eq (1) and -  
called particular integral.

As already pointed, the polynomial  $F(D)$ , can be subjected to algebraic operations, such as factorisation, resolutions, in partial fractions and expansions by binomial theorem etc. The following results are quite useful to find particular integrals.

**Results :-**

(i) If  $x$  is a function of  $x$  or a const, then  $\frac{1}{D} x = \int x dx$

sol<sup>n</sup>:- Let  $y = \frac{1}{D} x$  operating both sides  $D$

$$Dy = D \left( \frac{1}{D} x \right) = x$$

or  $\frac{dy}{dx} = x$ , Integrating both side w.r.t  $x$   $y = \int x dx$ ,  
hence the result.



(1) If  $x$  is a function of  $x$  or a constant, then  $\frac{1}{(D-m)}x$   
 $= e^{mx} \int x e^{-mx} dx$

Sol<sup>n</sup>: Let  $\frac{1}{(D-m)}x = y$

$$(D-m)y = (D-m) \left( \frac{1}{(D-m)}x \right) = x$$

or  $\frac{dy}{dx} - my = x$ , which is linear

$$I.F = e^{\int -m dx} = e^{-mx}$$

Hence sol<sup>n</sup> is  $y e^{-mx} = \int x e^{-mx} dx$

$$y = e^{mx} \int x e^{-mx} dx \quad \therefore \frac{1}{D-m} x = e^{mx} \int x e^{-mx} dx$$

This formula enables us to evaluate the particular integral  $\frac{1}{F(D)}x$ ,

where  $F(D) = (D-m_1)(D-m_2) \dots (D-m_n)$

$$\therefore \frac{1}{F(D)} x = \frac{1}{(D-m_1)(D-m_2) \dots (D-m_n)} x$$

Resolving R.H.S. into partial functions.

$$\frac{1}{F(D)} x = \left( \frac{A_1}{D-m_1} + \frac{A_2}{D-m_2} + \dots + \frac{A_n}{D-m_n} \right) x$$

$$= A_1 e^{m_1 x} \int x e^{-m_1 x} dx + A_2 e^{m_2 x} \int x e^{-m_2 x} dx + \dots + A_n e^{m_n x} \int x e^{-m_n x} dx$$

## Particular Integrals in some special cases :

Let us consider the differential eq<sup>n</sup>.

$$(D^n + P_1 D^{n-1} + P_2 D^{n-2} + \dots + P_n) y = X$$

where all  $P_i$ 's are constant.

The above eq<sup>n</sup> can be written as  $F(D) y = X$

$$\text{particular integral} = \frac{1}{F(D)} X$$

Case 1:

$$\text{when } X = e^{ax}$$

$$D(e^{ax}) = ae^{ax}$$

$$D^2(e^{ax}) = a^2 e^{ax}$$

$$\dots\dots\dots D^n(e^{ax}) = a^n e^{ax} \dots\dots\dots (i)$$

Multiplying in (i) by  $P_n, P_{n-1}, \dots, P_1$  and respectively and adding,

$$(D^n + P_1 D^{n-1} + \dots + P_{n-1} D + P_n) e^{ax} = (a^n + P_1 a^{n-1} + \dots + P_{n-1} a + P_n) e^{ax}$$

$$\Rightarrow F(D) e^{ax} = F(a) e^{ax}$$

Operating both sides of eq<sup>n</sup> (ii) by  $\frac{1}{F(D)}$

$$\frac{1}{F(D)} [F(D) e^{ax}] = \frac{1}{F(D)} [F(a) e^{ax}]$$

$$\text{or } \frac{e^{ax}}{F(a)} = \frac{1}{F(D)} e^{ax}$$

which gives the particular integral of  $e^{ax}$

In case  $X = K$  (a constant) then

$$\frac{1}{F(D)} K = K \frac{1}{F(D)} e^{0x} = \frac{K}{F(0)}, \text{ provided } F(0) \neq 0$$

Case of Failure :- If  $F(a) = 0$ , the above method fails and we proceed as under.

Since  $F(a) = 0$

$D = a$ , is a root of  $F(D) = 0 \dots \dots (i)$

$(D-a)$  is a factor of  $F(D)$ , suppose  $F(D) = (D-a)F'(D)$ ,

where  $F'(a) \neq 0$ .

$$\text{Then } \frac{1}{F(D)} e^{ax} = \frac{1}{D-a} \cdot \frac{1}{F'(D)} e^{ax} = \frac{1}{D-a} \cdot \frac{1}{F'(a)} e^{ax}$$

$$= \frac{1}{F'(a)} \cdot \frac{1}{(D-a)} e^{ax} = \frac{1}{F'(a)} e^{ax} \int e^{ax} \cdot e^{-ax} dx$$

$$= \frac{1}{F'(a)} e^{ax} \int dx = x \cdot \frac{1}{F'(a)} e^{ax}$$

$$\text{i.e. } \frac{1}{F(D)} e^{ax} = x \frac{1}{F'(a)} e^{ax} \dots \dots (2) \quad (F'(a) \neq 0)$$

$$\left[ \begin{array}{l} \therefore F'(D) = (D-a)F''(D) + 1 \cdot F(D) \\ \therefore F'(a) = 0 \cdot F''(a) + F(a) \end{array} \right]$$

If  $F'(a) = 0$ , then applying (2) again

$$\text{we get } \frac{1}{F(D)} e^{ax} = x^2 \frac{1}{F''(a)} e^{ax} \text{ provided } F''(a) \neq 0 \dots (3)$$

and so on,

In general if  $D = a$ , is a repeated root of  $F(D) = 0$ , say  $(n+1)$  times,

$$\text{we have } \frac{1}{F(D)} e^{ax} = \frac{x^n \cdot e^{ax}}{F^{(n)}(a)} \quad F^{(n)}(a) \neq 0$$

Rule 1 :-

To find the P.I.  $\frac{1}{F(D)} e^{ax}$ , replace  $D$  by  $a$ , provided  $F(a) \neq 0$ ,  
In case  $F(a) = 0$  multiply the numerator by  $x$  and  
differentiate the denominator w.r.t.  $D$  and apply the  
above rule, provided  $F'(a) \neq 0$  and so on.



## -: Partial Differential Equation :-

- \* The equation which contain one or more partial derivative is called partial differential equation. They must involve at least two independent variable and one dependent variable. Whenever we consider the case of two independent variable we shall usually take them to be  $x$  and  $y$  and  $z$  to be the dependent variable.
- \* The partial differential co-efficients  $\frac{\partial z}{\partial x}$  &  $\frac{\partial z}{\partial y}$  will be denoted by  $Q$  and  $P$  respectively.
- \* The second order partial derivatives  $\frac{\partial^2 z}{\partial x^2}$ ,  $\frac{\partial^2 z}{\partial x \partial y}$ ,  $\frac{\partial^2 z}{\partial y^2}$   $r, s, t$  respectively.
- \* The order of a partial derivative or partial differential equation is same as the order of the highest partial derivative in the equation. And it's degree is the degree of this derivative.

for example :-

$$\textcircled{i} \quad x \frac{\partial z}{\partial y} + y \frac{\partial z}{\partial x} = z$$

here order 1, degree 1

$$\textcircled{ii} \quad \frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0$$

here

- \* The partial differential eq<sup>n</sup> by the elimination of the form either function from a relation involving three of variables.

consider  $z$  to be a function of  $x$  and  $y$  defined by  $F(x, y, z, a, b) = 0$  ————  $\textcircled{1}$  in which  $a$  and  $b$  are to ~~are~~ constant differencing equation partially w.r.t  $x$ .

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial z} \times \frac{\partial z}{\partial x} = 0$$

$$= \frac{\partial F}{\partial x} + P \frac{\partial F}{\partial z} = 0 \quad \text{—————} \textcircled{2}$$

differentiating eqn ① partially w.r.t eqn ② w.r.t  $y$

$$\frac{\partial F}{\partial y} + \frac{\partial F}{\partial z} \times \frac{\partial z}{\partial y} = 0$$

$$\Rightarrow \frac{\partial F}{\partial y} + q \frac{\partial F}{\partial z} = 0 \quad \text{--- ③}$$

By means of the eqn ①, ②, ③ to constants  $a$  and  $b$  can be eliminated, this results in a partial differential eqn of order 1 in the form  $F(x, y, z, p, q) = 0$

## FIND THE SOLUTION OF $F(D)Y = E^{ax}$ :-

General solution of  $F(D)y = x$  (Non-homogeneous linear equation)

Consider the non-homogeneous linear differential eq<sup>n</sup> of  $n$ th order with co-efficients.

$$(a_0 D^n + a_1 D^{n-1} + a_2 D^{n-2} + \dots + a_{n-1} D + a_n) y = x$$

where,

$a_0, a_1, \dots, a_n$  one rational co-efficients

$$\text{i.e. } F(D)y = x \text{ ————— ①}$$

where,

$$F(D) = a_0 D^n + a_1 D^{n-1} + a_2 D^{n-2} + \dots + a_{n-1} D + a_n$$

$x$  is a function of  $x$ .

→ The complete integral of equation ① consists of two parts namely the complementary function (C.F) & a particular integral (P.I)

The particular integral of eq<sup>n</sup> ① is denoted by

$$P.I = \frac{1}{F(D)} \cdot x$$

Now suppose  $F(D) = D - a$

The P.I is given by  $y = \frac{1}{D-a} x$

$$\Rightarrow (D-a)y = x$$

$$\Rightarrow \frac{dy}{dx} - ay = x$$

which is a linear eq<sup>n</sup> in  $y$

Hence the solution is

$$y = \frac{1}{D-a} x = e^{ax} \int e^{-ax} x \, dx$$

$$\therefore \frac{1}{D-a} x = e^{ax} \int e^{-ax} x \, dx \text{ ————— ②}$$

If  $F(D) = a_0 D^2 + a_1 D + a_2$  i.e second degree,

then we can express

$$F(D) = (D-\alpha)(D-\beta)$$

So, P.I is given by  $\frac{\alpha}{(D-\alpha)(D-\beta)}$

Resolve  $\frac{1}{(D-\alpha)(D-\beta)}$  into partial fractions and then use the relation eqn (2) to evaluate the P.I.

$\therefore$  General solution is

$$y = C.F + P.I$$

To find P.I. when  $x$  is of the form  $x = e^{ax}$

$$\text{The P.I} = \frac{1}{F(D)} x = \frac{1}{F(D)} e^{ax}$$

we know that

$$D(e^{ax}) = ae^{ax}$$

$$D^2(e^{ax}) = a^2 e^{ax}$$

$\vdots$

$$D^n(e^{ax}) = a^n e^{ax}$$

$$\therefore F(D) e^{ax} = F(a) e^{ax}$$

$$\frac{1}{F(D)} [F(D) e^{ax}] = \frac{1}{F(D)} [F(a) e^{ax}] = F(a) \frac{1}{F(D)} e^{ax}$$

$$\frac{1}{F(D)} [F(D) e^{ax}] = F(a) \frac{1}{F(D)} e^{ax}$$

$$\Rightarrow e^{ax} = F(a) \cdot \frac{e^{ax}}{F(D)}$$

$$\Rightarrow \frac{1}{F(D)} e^{ax} = \frac{e^{ax}}{F(a)} \quad \text{provided } F(a) \neq 0$$

Now if

$F(a) = 0$ , then  $F(D) = (D-a)g(D)$  where  $g(a) \neq 0$



So,

$$PI = \frac{1}{F(D)} x$$

$$= \frac{1}{F(D)} \cdot e^{ax} = \frac{e^{ax}}{(D-a)g(D)} = \frac{1}{g(a)} \frac{e^{ax}}{D-a}$$

$$= \frac{1}{g(a)} \left[ e^{ax} \int e^{-ax} \cdot e^{ax} dx \right]$$

$$\left( \because \frac{1}{D-a} = e^{ax} \int e^{-ax} x dx \right)$$

$$\Rightarrow P.I = \frac{1}{g(a)} \left[ e^{ax} \int dx \right] = \frac{x e^{ax}}{g(a)}$$

we can also take

$g(D) = F(D)$  here

Similarly,

$$\frac{e^{ax}}{(D-a)^2} = \frac{x^2 e^{ax}}{2!} \quad \text{and} \quad \frac{e^{ax}}{(D-a)^3} = \frac{x^3 e^{ax}}{3!}$$

Working rule for evaluating P.I for  $F(D)y = e^{ax}$  :-

Write the factors of  $\psi(D) = (D-a)(D-b)$

Substitute  $D = a$  if  $F(a) \neq 0$

If  $F(a) = 0$ , then

$$\frac{1}{F(D)} e^{ax} = \frac{1}{(D-a)} e^{ax} = \frac{x}{1} e^{ax},$$

$$\frac{1}{(D-a)^2} e^{ax} = \frac{x^2}{2!} e^{ax}$$

$$\frac{1}{(D-a)^3} e^{ax} = \frac{x^3}{3!} e^{ax}$$

Ex 1 solve  $(D^2 - 7D + 6)y = e^{2x}$

Ex 2 solve  $\frac{d^2 y}{dx^2} + \frac{dy}{dx} + 3y = e^{2x}$

3. Solve  $\frac{d^2y}{dx^2} - 5 \frac{dy}{dx} + 6y = e^{4x}$

4. Solve  $(4D^2 + 4D - 3)y = e^{2x}$

5. Solve  $\frac{d^2y}{dx^2} + \frac{dy}{dx} + y = e^{4x}$

6. Solve  $\frac{d^2y}{dx^2} + 5 \frac{dy}{dx} + 6y = e^{-2x}$

7. Solve  $(D^2 + 6D + 5)y = 16e^{3x} + 2e^{-x} + 3$

8. Solve  $y'' + 4y' + 5y = -2 \cosh x$ . Also find  $y$  where  $y=0$ ,  $\frac{dy}{dx} = 1$  at  $x=0$

Find the solution of  $f(D)y = \cos ax$  or  $\sin ax$ . :-

$\Rightarrow$  Particular Integral of  $F(D) = x$  when  $x = \cos ax$  or  $\sin ax$  where 'a' is any constant.

Let  $x = \sin ax$  then

$$P.I = \frac{1}{F(D)} x = \frac{1}{F(D)} \sin ax$$

we have.

$$D(\sin ax) = a \cos ax, D^2(\sin ax) = -a^2 \sin ax$$

$$D^3(\sin ax) = -a^3 \cos ax, D^4(\sin ax) = a^4 \sin ax$$

$$D^5(\sin ax) = a^5 \cos ax, D^6(\sin ax) = -a^6 \sin ax \text{ and so on}$$

Hence from the above it is clear that

$$D^2(\sin ax) = -a^2 \sin ax$$

$$D^4(\sin ax) = a^4 \sin ax = (-a^2)^2 \sin ax$$

$$D^6(\sin ax) = (-a^2)^3 \sin ax = -a^6 \sin ax$$

$$D^{2n}(\sin ax) = (D^2)^n \sin ax = (-a^2)^n \sin ax$$

$$\text{Since } F(D) = D^n + a_1 D^{n-1} - a_2 D^{n-2} + \dots + a_n$$

$$\therefore F(D^2) \sin ax = \left[ (D^2)^n + a_1 (D^2)^{n-1} + a_2 (D^2)^{n-2} + \dots + a_n \right] \sin ax$$

$$= \left[ (-a^2)^n \sin ax + a_1 (-a^2)^{n-1} \sin ax + a_2 (-a^2)^{n-2} \sin ax + \dots + a_n \sin ax \right]$$

$$= \left[ (-a^2)^n + a_1 (-a^2)^{n-1} + a_2 (-a^2)^{n-2} + \dots + a_n \right] \sin ax$$

$$\Rightarrow F(D^2) \sin ax = F(-a^2) \sin ax$$

Operating on both sides by  $\frac{1}{F(D^2)}$ , we get

$$\frac{F(D^2) \sin ax}{F(D^2)} = \frac{F(-a^2) \sin ax}{F(D^2)} \quad \therefore F(-a^2) \text{ is a const.}$$

$$\Rightarrow \sin ax = F(-a^2) \frac{1}{F(D^2)} \sin ax$$

$$\frac{F(D^2) \sin ax}{F(D^2)} = \frac{F(-a^2) \sin ax}{F(D^2)} \quad \therefore F(-a^2) \text{ is a const.}$$

$$\Rightarrow \sin ax = F(-a^2) \frac{1}{F(D^2)} \sin ax$$

$$\Rightarrow \frac{1}{F(D^2)} \sin(ax) \frac{1}{F(-a^2)} \sin ax, \text{ if } F(-a^2) \neq 0$$

$$\text{Hence P.I} = \frac{1}{F(D^2)} \sin ax, \text{ if } F(-a^2) \neq 0$$

$$= \frac{1}{F(-a^2)} \sin$$